



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

PROBLEMS FOR SOLUTION.

[N.B. The editorial work of this department would be greatly facilitated if, on sending in problems, the proposers would also enclose their solutions—*when they have them*. If a problem proposed is not original the proposer is requested *invariably* to state the fact and to give an exact reference to the source.]

2890. Proposed by B. F. FINKEL, Drury College.

Having given a triangle whose base is $2c$ and (a) the sum of whose other two sides is $2a$, (b) the difference of whose other two sides is $2a$, determine the envelope of the perpendicular bisectors of the variable sides.

2891. Proposed by D. F. BARROW, Philomath, Ga.

Let A' , A'' , A''' , and P denote, respectively, the vertices of a triangle and any point in its plane; and let P' , P'' , P''' denote the feet of the perpendiculars from P upon the sides opposite A' , A'' , A''' . Now suppose each of the lines PP' , PP'' , PP''' to revolve about P through an angle α ; and let P_a' , P_a'' , P_a''' denote the intersections of this new triad of lines with the corresponding sides of the triangle. As α varies, find the envelope of the variable circle through P_a' , P_a'' , P_a''' .

2892. Proposed by R. T. MCGREGOR, Bangor, Calif.

Two parabolas have parallel axes. Prove that their common chord bisects their common tangent.

2893. Proposed by NATHAN ALTSHILLER-COURT, University of Oklahoma.

Find the locus of the mid-point of the segment determined by two given skew lines in a variable plane turning about a fixed axis, not coplanar with either of the given lines.

2894. Proposed by PHILIP FRANKLIN AND E. L. POST, Princeton University.

Given the following set of assumptions concerning a set S and certain undefined sub-classes of S , called m -classes:

I. If A and B are distinct elements of S , there is at least one m -class containing both A and B .

II. If A and B are distinct elements of S , there is not more than one m -class containing both A and B .

Def. Two m -classes with no elements in common are called *conjugates*.

III. For every m -class there is at least one *conjugate* m -class.

IV. For every m -class there is not more than one *conjugate* m -class.

V. There exists at least one m -class.

VI. Every m -class contains at least one element of S .

VII. Every m -class contains not more than a finite number of elements.

Develop some of the propositions of the "mathematical science" (cf. Veblen and Young, *Projective Geometry*, Vol. I, pp. 1 f.) based on them and in particular develop a sufficient number of theorems to prove that the set of assumptions is categorical and give a concrete representation of the set S which satisfies them. Also prove that the assumptions are independent.

2895. Proposed by R. M. MATHEWS, Wesleyan University.

To construct an equilateral triangle with its vertices lying on: (a) any three coplanar lines; (b) three parallels in space; and (c) any three lines in space.

PROBLEMS—NOTES

10. A Curve of Pursuit. The extended discussion of a curve of pursuit in a recent issue of this MONTHLY (1921, 54–61, 91–97) suggests this note. In *Nouvelle Correspondance Mathématique*, volume 3, 1877, E. Lucas proposed the following problem in May (pages 175–176): "Three dogs are placed at the vertices of an equilateral triangle; they run one after the other. What is the curve described